

Non-Clinical Value of ENF-VT3

Designed for Office-Based Laryngology



ENF-VT3

Designed for Office-Based Laryngology







· ENF-VT3 enables the move of procedures like biopsies to an outpatient setting, which may result in significant cost reduction of up to 91% per procedure.¹

- · Performing biopsies in an outpatient setting with ENF-VT3 could improve patient pathway by decreasing patients' waiting times by up to 93%.^{2,3}
- The unique four-direction angulation of the ENF-VT3 scope helps specialist to access difficult areas easier and with more precision.
- NBI* may further improve the diagnostic outcome of the biopsy by improving the contrast between suspicious lesions and healthy tissue.

* Narrow Band Imaging

Value Dimensions

ENF-VT3 provides a number of values for you and your staff

Safety **Patient Well-Being**

- Reduced waiting times may improve clinical outcome.
- Patients may be able to



Economics Cost and Cash-Flow Management

• Outpatient setting reduces cost per procedure and frees up hospital beds.

avoid hospital admission

altogether.

• Operating room capacity is

realeased for more complex

and highly reimbursed

procedures.

ENF-VT3

Designed for Office-Based Laryngology

Outpatient Biopsies Improve Patient Pathway by Reducing Waiting Times Significantly

Traditionally, patients with suspicious head and neck lesions would have a biopsy taken in an operating theater under general anesthesia to confirm a diagnosis. Challenges with scheduling, availability of theatre time, hospital beds and pre-procedure assessments can delay diagnosis and, where needed, the initiation of treatment.

Shifting to outpatient biopsy using ENF-VT3, could reduce the wait for a confirmed diagnosis and the time to start treatment. This approach has the potential to reduce the time between consultation and diagnosis by an average of more than two weeks (15.5 days) according to Lee at al. (2018). and may reduce the time from consultation to initiation of treatment by 14.5 days according to Shutte et al. (2018). Embracing this change is crucial to improving outcomes and addressing head and neck cancer promptly.

Average Waiting Time in Days from Consultation to Initiation of Treatment



UK NHS Example of Cost Savings When Moving from OR to Outpatient Setting



Cost per Procedure

** Since fewer staff are required in the outpatient setting, general anesthesia is avoided, and hospital bed stay is not required.

ENF-VT3

References

- Outpatient Procedures vs Day Case, CA69A Diagnostic, Laryngoscopy or Pharyngoscopy, 19 years and over, National Schedule of NHS Costs Year: 2020-21 — All NHS trusts and NHS foundation trusts — HRG Data, NHS England, 2022
- Decrease time from first consultation to biopsy 17.4 days (OR) vs 1.3 days (outpatient), and decrease time from first consultation to diagnosis — 23 days (OR) vs 7.5 days (outpatient): Lee F, Smith KA, Chandarana S, et al. (2018). An evaluation of in-office flexible fiber-optic biopsies for laryngopharyngeal lesions. Journal of Otolaryngology: Head & Neck Surgery. 47(1): 31. doi: https://dx.doi.org/10.1186/ s40463-018-0275-x
- Decrease time from first consultation to treatment 41.5 days (OR) vs 27 days (outpatient): Schutte HW, Takes RP, Slootweg PJ, et al. (2018). Digital video laryngoscopy and flexible endoscopic biopsies as an alternative diagnostic workup in laryngopharyngeal cancer: a prospective clinical study. Annals of Otology, Rhinology & Laryngology. 127(11): 770-6. doi: https://doi.org/10.1177/0003489418793987

S00603EN · 11/23 · OKM

Read More About ...

As medical knowledge is constantly growing, technical modifications or changes to the product design, product specifications, accessories and service offerings may be required.



OLYMPUS EUROPA SE & CO. KG

Postbox 10 49 08, 20034 Hamburg, Germany

Wendenstrasse 20, 20097 Hamburg, Germany

Phone: +49 40 23773-0, Fax: +49 40 233765

www.olympus-europa.com